

## **Lower Body Orthosis** Custom Measurement Form

					Date	
Name (Patient)					Age	
Contact Name						
Contact Phone		Contact E	Email			
Measurement Key	ALL	. MEASUR	EMENT	FIELDS AF	RE REQUIR	ED
Length =					and the same	Vaist
Circumference =			A	. G	(all	navel)
Instructions					/ TI	rochanter Line
-Measure patient	RIGHT		В	H <b>∠</b>	LĘ	FT
standing upMeasurements should	Waist line (at navel) to Trochanter	Top of Thigh		7	Top of Thigh	Waist line (at navel) to Trochanter
be in centimeters.	cm	(cm)			cm)	cm
-Measure knee circumference at 15	Line A - B		*			Line G - H
degrees flexion.	Waist line (at navel)	Above Knee Cap (at Condyle)	\		Above Knee Cap (at Condyle)	Waist line (at navel)
-Measure both legs.	to Condyle		_ \ c			to Condyle
-Measure over diaper or	cm		<b>-</b>		cm	cm
underwear.	Line A - C	Mid Knee Cap	D ]	<u>\                                    </u>	Mid Knee Cap	Line G - I
-All measurements must be provided.	Waist line (at navel) to Mid Knee		×     /	1		Waist line (at navel) to Mid Knee
-Measure on top of torso,	to wild knee	cm)			cm	- I I I I I I I I I I I I I I I I I I I
not the side.	cm	Calf (fullest part)	1   1		Calf (fullest part)	cm
	Line A - D	Carr (rulest part)	E .	K	Can (rulest part)	Line G - J
	Waist line (at navel)	(cm)-	7	1 1	—(cm)	Waist line (at navel)
Diaper	to Calf (fullest part)		1   (	) /		to Calf (fullest part)
OYES ONO	cm	Ankle	\ \		Ankle	cm
0	Line A - E	(cm)_	F		_(cm)	Line G - K
If yes, measure back waist	Waist line (at navel) to Ankle			1 1 -		Waist line (at navel) to Ankle
to gluteal foldcm	WAIRIE		1 1			to Ankie
	cm		(,,,)	1 )		cm
Overstitus	Line A - F			COM		Line G - L
Quantity	Inseam (crotch to hemlir	ne)				Inseam (crotch to hemline
	cm					cm
	Line M - C, D, E, or F					Line M - I, J, K, or L
Options			Additio	onal Comments	<b>Š</b> @	
	ROYAL BLUE	WHIT	·c			
		_				
Hem ( )CONDYLE (	MID KNEE	CALF ( ) AN	KLE			